

UVES observations of QSO 0000–2620: oxygen and zinc abundances in the Damped Ly α galaxy at $z_{abs} = 3.3901$ ¹

Paolo Molaro¹, Piercarlo Bonifacio¹, Miriam Centurión¹, Sandro D’Odorico², Giovanni Vladilo¹, Paolo Santin¹ and Paolo Di Marcantonio¹

¹ Osservatorio Astronomico di Trieste, Via G.B. Tiepolo 11, 34131, Trieste

²European Southern Observatory, Karl Schwarzschildstr. 2, D-85740 Garching, Germany

Received _____; accepted _____

¹Based on observations made with the ESO 8.2m KUEYEN telescope operated on Paranal Observatory, Chile.

ABSTRACT

Observations of the QSO 0000–2620 with UVES spectrograph at the 8.2m ESO KUEYEN telescope are used for abundance analysis of the damped Ly α system at $z_{abs}=3.3901$. Several Oxygen lines are identified in the Ly α forest and a measure for the oxygen abundance is obtained at $[O/H]=-1.85 \pm 0.1$ by means of the unsaturated OI λ 925Å and OI λ 950Å lines. This represents the most accurate O measurement in a damped Ly α galaxy so far. We have also detected ZnII λ 2026 Å and CrII λ 2056, λ 2062 Å redshifted at ≈ 8900 Å and found abundances $[Zn/H] = -2.07 \pm 0.1$ and $[Cr/H]=-1.99 \pm 0.1$. Furthermore, previous measurements of Fe, Si, Ni and N have been refined yielding $[Fe/H]=-2.04 \pm 0.1$, $[Si/H]=-1.90 \pm 0.1$, $[Ni/H]=-2.27 \pm 0.1$, and $[N/H]=-2.68 \pm 0.1$. The abundance of the non-refractory element zinc is the lowest among the damped Ly α systems showing that the associated intervening galaxy is indeed in the early stages of its chemical evolution. The fact that the Zn abundance is identical to that of the refractory elements Fe and Cr suggests that dust grains have not formed yet.

In this Damped Ly α system the observed [O,S,Si/Zn,Fe,Cr] ratios, in whatever combination are taken, are close to solar (i.e 0.1–0.2 dex) and do not show the $[\alpha\text{-element}/Fe]$ enhancement observed in Milky Way stars of comparable metallicity. The observed behavior supports a galaxy evolution model characterized by either episodic or low star formation rate rather than a Milky-Way-type evolutionary model.

Subject headings: cosmology: observations — galaxies: abundances — galaxies: evolution — quasars: absorption lines

1. Introduction

Damped Ly α (DLA) systems ($\log N(\text{HI}) \geq 20.3 \text{ cm}^{-2}$) are seen in absorption in the QSO spectra up to the highest redshift. They likely trace the field population of galaxies with no a priori selection based on morphology, since biases apply to the background sources at first approximation and not to the intervening systems. The identification of the galaxy population associated with the DLAs has important bearings on cosmological elemental evolution and has been actively debated.

Wolfe et al. (1995) and Prochaska & Wolfe (1997,1999) suggested a close connection between the DLA galaxies and the progenitors of present-day spiral galaxies. The connection is based on the fact that the comoving gas density of the DLA galaxies at high redshift roughly equals the local density provided by the luminous matter in nearby spiral galaxies. In addition, the kinematics of the metal lines has been interpreted as having formed in massive rotating disks. However, Rao & Turnshek (2000) did not find a decrease in the comoving gas density of DLA galaxies at low redshifts as expected if the gas had been converted into stars, and Haehnelt, Steinmetz, & Rauch (1998) and Ledoux et al. (1998) did not find the kinematical argument very compelling.

DLA systems offer the possibility to measure abundances accurately for a wide range of elements in a variety of normal, gas-rich galaxies spanning a considerable look-back time. The metallicity of DLA galaxies is generally inferred from the absolute abundance of the volatile element Zinc which is found at $\sim 10\%$ of solar, with no clear evolution over the redshift interval from $z_{abs} \approx 0.5$ up to 3 (Pettini et al. 1999). The apparent lack of evolution has been considered a problem for the interpretation of DLAs as proto-spirals, in particular at low redshift where abundances much closer to the solar ones are expected. The models of Fritze et al. (1999) for Sa and Sd galaxies bracket the Zn data from the high and low metallicity sides over the entire redshift range, but require that DLAs are biased against early type spirals at low redshift. A possible bias is represented by the low gas and the high dust contents of early type galaxies at low redshift. On the same line, Prantzos and Boisser (2000) found the observed Zn abundances consistent with a spiral origin once the empirical relation $18.8 \leq [\text{Zn}/\text{H}] + \log N(\text{HI}) \leq 21$ is imposed to the models. Jimenez et al. (1999) found that the population of Low Surface Brighness galaxies (LSB) better matches the observed zinc metallicities. Imaging of DLA galaxies at relatively low redshift revealed that the population of DLA galaxies is not dominated by spirals but rather composed by a variety of morphological types which include LSB, dwarf irregulars and late type spirals. (Le Brun et al. 1997; Rao & Turnshek 1998).

Relative elemental abundances are an important complement to the information provided by absolute

abundances such as zinc metallicity. In particular, ratios of elements which are produced by Type I and Type II supernovae in different proportions provide independent insights into defining the kind of chemical evolution. Despite the efforts of different authors, it has been difficult to establish the $[\alpha/\text{Fe}]$ abundance pattern in DLA systems because of the different level of depletion of refractory elements onto dust grains. In the compilation of Savaglio et al. (2000) the average of 37 measures is $\langle [\text{Si}/\text{Fe}] \rangle = +0.43 \pm 0.18$ ². At face value this is consistent with what is observed in the Galactic halo stars, which has been interpreted as evidence for an origin of DLAs in proto-spirals (Lu et al. 1996, Prochaska & Wolfe 1999).

Evidence that at least in some DLAs relative abundances do not conform to those of the Galactic halo stars was first provided by Molaro et al. (1996) in a first study of the abundances in the DLA at $z_{\text{abs}}=3.3901$ towards QSO 0000–2620, which is considered again in this paper. Molaro, Centuri3n, & Vladilo (1998) and Centuri3n et al. (2000) also found the ratio of the undepleted elements Sulphur and Zinc to be approximately solar in a sample of half a dozen DLAs. Vladilo (1998) corrected the abundances of Si and Fe for the differential elemental depletion and obtained approximate solar ratios ($[\text{Si}/\text{Fe}] \approx 0$) in all cases investigated, thus suggesting that solar ratios might be rather common among DLAs. Pettini et al. (2000) considering DLAs with low dust depletion found $[\text{Si}/\text{Fe}]$ ratios broadly in line with Galactic stellar values although there are also examples of near-solar $[\text{Si}/\text{Fe}]$ ratios at $[\text{Fe}/\text{H}] < -1$. Savaglio et al. (2000), after correcting for dust, found evidence for overabundance of a factor 2 in Si in about 1/4 of the DLA systems.

The $[\text{O}/\text{Zn}]$ ratio is probably the best diagnostic tool for the $[\alpha/\text{Fe}]$ ratio we have for DLAs since Oxygen is produced by TypeII SNe and is essentially undepleted in the interstellar medium. Unfortunately, it is difficult to obtain the oxygen abundance with the required accuracy. In the present study of the $z=3.3901$ DLA system towards QSO 0000–2620 we present an accurate oxygen abundance determination based on unsaturated lines, together with a Zn detection at the highest redshift ever obtained, and discuss the $[\text{O}/\text{Zn}]$ ratio in connection with the chemical evolution of the associated galaxy.

2. Observations and data reduction

QSO 0000–2620 is a bright QSO ($V=17.5$) with $z_{\text{em}}=4.108$ discovered by C. Hazard, which shows a damped $\text{Ly}\alpha$ system at $z_{\text{abs}} = 3.3901$. This is one of the few damped systems at redshift greater than 3 known so far and is well suited for a detailed study of chemical abundances in primordial galaxies.

²Using the customary definition $[\text{X}/\text{H}] = \log (\text{X}/\text{HI}) - \log (\text{X}/\text{H})_{\odot}$

Measurements of abundances of this system have been obtained by Savaglio et al. (1994) and Molaro et al. (1996) from EMMI observations at the ESO NTT, and by Lu et al. (1996) and Prochaska & Wolfe (1999) from HIRES observations at KeckI.

A few spectra of QSO 0000–2620 were obtained as test observations during the first commissioning of the Ultraviolet-Visual Echelle Spectrograph (UVES) on the Nasmyth focus of the ESO 8.2m KUEYEN telescope at Paranal, Chile, in October 1999, which were released for public use. In this paper we have used the data of higher quality: two exposures of 4000 sec and one of 4500 sec secured on October 13, when the seeing, as given by the telescope guide probe, was between 0.35 and 0.5 arcsec FWHM. The observations were all obtained in the standard dichroic #2 mode which includes the spectral region from 3700 to 5000 Å in the blue arm and from 6700 Å to 10500 Å in the red arm. The CCD in the blue arm is a 2Kx4K, 15 μm pixel size thinned, anti-reflection coated EEV CCD-44, while in the red the detector is a mosaic of an EEV CCD-44 of the same type of the blue arm and a MIT/LL CCID-20. The average pixel scale in the direction of the dispersion is 0.22 and 0.16 arcsec/pixel for the blue and red arm, respectively. The slit width was set at 0.9 arcsec. The CCDs were read-out in 2x2 pixel binned mode, which gives a r.o.n. of 2.1 e[−] rms for the blue arm CCD, and 2.1 and 3.4 e[−] rms for the two red arm CCDs. More details about the instrument can be found in Dekker, D’Odorico and Kaufer (2000). The full width at half maximum of the instrumental profile, $\Delta\lambda_{instr}$, was measured from the emission lines of the Thorium-Argon lamp. The resulting resolving power is $R = \lambda/\Delta\lambda_{instr} \simeq 48000$, which corresponds to a velocity resolution of $\simeq 6 \text{ km s}^{-1}$.

The data reduction was performed using the ECHELLE context routines implemented in the ESO MIDAS package. Flat-fielding, cosmic ray removal, sky subtraction, and wavelength calibration were performed on each spectrum separately. Typical r.m.s of the wavelength calibrations are $\leq 0.6 \text{ km s}^{-1}$.

The observed wavelength scale of each spectrum was then transformed into vacuum, heliocentric wavelength scale. The spectra were then added together by using their S/N as weights. Finally, the local continuum was determined in the average spectrum by using a spline to smoothly connect the regions free from absorption features. The continuum for the Ly α forest region was fitted by using the small regions deemed to be free of absorptions and by interpolating between these regions with a spline.

With respect to the previous ESO NTT observations, UVES spectra have a higher extension into the blue as well as a more than double resolution and higher S/N. With respect to the Keck data, which cover the range 5100–7660 Å, UVES spectra extend further towards both the blue and the red sides. In the region of overlap, which extends from ~ 6700 to 7660 Å the data have similar resolution and comparable S/N as

can be seen from a comparison of our figures with Fig.1 of Prochaska & Wolfe (1999).

3. Measurements of the Column Densities

In this work we restricted our analysis to the metal lines falling in the Ly α forest which have never been observed at both comparable resolution and S/N, and to lines in the reddest part of the spectrum. Measurements are presented in Table 1. Column densities have been obtained by fitting theoretical Voigt profiles to the observed absorption lines via χ^2 minimization. The fit was performed using the FITLYMAN package within MIDAS (Fontana & Ballester 1995). During the fitting procedure the theoretical profiles were convolved with the instrumental point spread function modeled from the analysis of the emission lines of the arcs. Portions of the profiles recognized as contaminated by intervening Ly α clouds were excluded from the analysis.

The FITLYMAN routines determine the redshift, the column density, and the broadening parameter (b -value)³ of the absorption components, as well as the fit errors for each quantity. The atomic data were taken from the compilation of Morton (1991) with the revisions by Cardelli & Savage (1995), Bergeson & Lawler (1993a,b) and Fedchak and Lawler (1999), as specified in Table 1.

Since Ly α is not included in our range, when calculating the abundances the hydrogen column density is taken as $\log N(HI)=21.41\pm0.08$ from Lu et al. (1996), after having verified the value consistent with the Ly β absorption. With such a large column density, the ion species are from the dominant ionization stages in HI gas and do not require ionization corrections (Viegas 1995). The damped structure is known to include minor additional components on both sides of the main absorption (Molaro et al. 1996, Lu et al. 1996), but our analysis is restricted to the unsaturated lines where these weak components do not appear and do not affect the abundances.

The derived column densities and b -values, together with the redshifts, are reported in Table 1. The difficulty in determining accurate OI column densities stems from the fact that the only transition available redwards the Ly α emission is OI λ 1302Å which has always been found heavily saturated in DLAs. On the other hand OI λ 1356 Å has never been detected and has provided only loose upper limits. Upon searching

³The broadening parameter is defined as $b = 2^{1/2}\sigma_v$, where σ_v is the one-dimensional gaussian velocity dispersion of ions along the line of sight.

the Ly α forest we detected several oxygen lines, namely OI λ 925.446, λ 929.517, λ 936.629, λ 948.685, λ 950.884, λ 988.773, λ 1039.230 Å which are relatively free from contamination caused by hydrogen clouds. The OI λ 925 Å and OI λ 950 Å lines have a very low f value (3.5×10^{-4} and 15.7×10^{-4}), which is 140 and 25 times lower than the OI λ 1302 Å; respectively. The lines are not saturated thus allowing accurate determination of the oxygen abundance. The two oxygen lines are shown in Fig. 1. Owing to the presence of minor contaminants we analyzed the two lines independently. The two lines yield similar column densities, where that obtained from OI λ 925 Å is slightly higher. We notice that the OI λ 925 Å is somewhat broader ($b=14$ km s $^{-1}$) than the other metal lines, thus suggesting the presence of some blend which may explain the small difference in the column densities. In fitting OI λ 950 the presence of the wing of the Ly δ line on the blue side is accounted for. The adopted abundance for OI is taken as the straight average of the two measurements, i.e. $\log N(\text{OI})=16.42$ (or $[\text{O}/\text{H}]=-1.85 \pm 0.1$) with an error of ± 0.1 which accounts for the uncertainty in the continuum placement. The other α -element, Silicon, is obtained from the unsaturated SiII λ 1808 Å line also shown in Fig. 1. We obtained a column density of $\log N(\text{SiII})=15.06$, which is in good agreement with the value of 15.086 derived by Prochaska & Wolfe (1999).

Several N transitions belonging to the NI λ 952, λ 953, λ 956, λ 1134 Å multiplets have been detected for the first time in a DLA. In each multiplet, however, only few lines resulted relatively free from contaminations, namely NI λ 952.415, λ 953.415, λ 953.655, NI λ 963.990, and λ 965.041 Å. In deriving the nitrogen abundance we further restricted the analysis to those lines which appear totally free from contaminations, which comprise the three lines of the NI λ 1134 Å multiplet and the NI λ 953.6 Å transition and are shown in Fig 2. The N column density is found $\log (NI) = 14.70$, which is in excellent agreement with the $\log N(\text{NI})= 14.68 \pm 0.14$ derived by Molaro et al. (1996) from the analysis of the NI λ 1134 Å multiplet.

The iron abundance is obtained from the unsaturated FeI λ 1611 Å line. The abundance is determined at $[\text{Fe}/\text{H}]=-2.04$, which is 0.3 dex higher than the value obtained by Molaro et al. (1996), and 0.2 dex lower than that obtained by Prochaska and Wolfe (1999) by means of the same transition. The FeII line together with the other iron-peak elements are shown in Fig. 3.

Chromium is clearly detected through the CrII λ 2056 Å and λ 2062 Å transitions, while the CrII λ 2066 Å line is obscured by a strong sky emission line. The two lines yield almost identical Cr column densities with a mean value of $\log N(\text{CrII})=13.10$.

Zinc abundance is obtained from the stronger component of the doublet Zn II λ 2026.1360 Å while the

weaker ZnII λ 2062.664 Å remains below the detection threshold. The value found here is $[\text{Zn}/\text{H}]=-2.07$ which is consistent with the upper limits set at $[\text{Zn}/\text{H}]\leq -1.76$ (3σ) by Pettini et al. (1995).

We have clearly detected the NiII λ 1709 Å while the NiII λ 1751 Å showed to be partially blended with an O₂ atmospheric line. The abundances of NiII λ 1709 Å yields $[\text{Ni}/\text{H}]=-2.27$, which is also consistent with the value of NiII λ 1751 Å when the telluric contamination is accounted for. We note that the abundance of Ni, despite the recent upwards revision in the f values by Fedchak and Lawler (1999), remains slightly lower than the other iron-peak elements.

Other transitions present in our spectrum such as AlIII and CIV are either saturated or are not significantly different from the values published in the quoted references. In this paper they are not examined again.

4. Discussion

4.1. Zinc Abundance and Dust Content

Zinc abundances have been used to trace back the cosmological chemical evolution of Damped Ly α galaxies. Instead of the expected growth of metallicity with cosmic time, which is a natural outcome of stellar evolution, Pettini et al. (1999) could not find any trend of the $[\text{Zn}/\text{H}]$ ratio with z_{abs} . The DLA towards QSO 0000–2620 is the only one at $z \geq 3$ for which a Zn abundance measurement is available at present. The zinc abundance is found at $[\text{Zn}/\text{H}]=-2.07$, which is the lowest level found so far in any DLA at any redshift. The low metallicity level shows that this Ly α galaxy is indeed in the early stages of its chemical evolution. When compared with extant data, this Zn abundance suggests that a mild chemical evolution is present in the DLA population. Observational biases may affect this conclusion, as discussed in a separate paper (Vladilo et al. 2000).

A second important point is that zinc abundance at $[\text{Zn}/\text{H}]=-2.07$ is fully consistent with both iron and chromium abundances, which are $[\text{Fe}/\text{H}]=-2.04$ and $[\text{Cr}/\text{H}]=-2.0$ respectively. It is well established that in the DLAs Zn is consistently more abundant than Cr and Fe (Pettini et al. 1994, 1997), while in Galactic halo stars both Cr and Zn track the Fe content down to very low metallicities. The different behavior of Fe and Zn in the DLAs is currently interpreted as the result of differential depletion of these elements from the gas phase onto dust grains in analogy with the nearby interstellar medium. The fact that in this DLA the abundances of the refractory elements Fe and Cr are found close to the non-refractory element Zn suggests

that dust grains have not yet formed in this protogalaxy. Such a circumstance is extremely rare among DLAs, and even in the cases with low depletion considered by Pettini et al. (2000), a difference between Zn and the other refractory elements Fe, Cr and Ni has been always observed. The absence of dust in this damped system frees the measurements of relative abundances from the uncertainties deriving from the unknown fraction of elements which condensate onto grains, thus making this DLA of particular value in the study of chemical composition patterns.

4.2. Abundances of α -capture versus iron-peak elements

Information on α -capture elements is important because in the early stages of the chemical evolution of galaxies the abundances are likely dominated by Type II SNe products which are richer in α -elements than Type I SNe which entered into the game only later on. Thus in the early stages the α -capture elements are expected to be relatively more abundant than the iron-peak elements, and the ratio should reverse later on during evolution. The exact timing of the turn-over depends on both the SFR and IMF, which differ for the different types of galaxies. The $[\alpha/\text{Fe}]$ ratio is therefore an indicator of the chemical evolution history which can be used to understand the nature of galaxies when data on their morphology and colors are lacking. Silicon is the most accessible α element in DLAs and the compilation of Savaglio et al. (2000), who consider 37 measures, obtain a mean value of $\langle [\text{Si}/\text{Fe}] \rangle = 0.43 \pm 0.18$. The problem here is that we observe a similar ratio ($[\text{Si}/\text{Fe}] = +0.36$) in the Warm Phase of the Galactic interstellar medium, due to the fact that the gaseous silicon and iron abundances are lowered by different amounts by dust depletion (Savage & Sembach 1996). After accounting for dust, Savaglio et al. (2000) conclude that Si is intrinsically enhanced in comparison to iron only in about 25% of the DLAs with SiII determined. Outram et al. (1999) in two DLA systems inferred the absence of dust from the similarity of Si and S abundances and argued for an intrinsic overabundance of $[\alpha/\text{Fe}]$ of about 0.5 dex. However, as noted in Matteucci et al. (1997), Si might be produced in Type I SNe, which could explain the complicate observational pattern. Therefore, Si should be considered an indicator of α element abundances less reliable than O or S.

Evidence that relative abundances in some DLAs do not conform to those of the galactic halo stars has been provided by the $[\text{S}/\text{Zn}]$ ratio (Molaro et al. 1996, 1998, Centuri3n et al. 2000) in a sample of 6 DLAs but with one case, the DLA at $z_{\text{abs}} = 2.476$ towards Q0841+129, consistent with $[\text{S}/\text{Zn}] > 0.2$.

The oxygen abundance we derive here is the first one obtained by means of unsaturated lines. Since the transitions lie in the forest, the uncertainty is dominated by possible $\text{Ly}\alpha$ interlopers. However, if these

hydrogen lines do actually contaminate the spectrum at the wavelength of the oxygen lines, the real oxygen abundance should be even lower than the value found here.

Oxygen is a typical product of type II SNe and in the atmospheres of Galactic metal-poor stars it shows the notorious enhancement of $[O/Fe] \simeq +0.5$. Claims for an even more extreme oxygen overabundance with $[O/Fe]$ reaching +1.0 dex at $[Fe/H]=-3.0$ have been made recently by Israelian et al. (1998) and Boesgaard et al. (1999).

Oxygen has a rather small condensation temperature ($T_c \approx 180$ K) and its depletion factor is of $0.00^{+0.18}_{-0.31}$ dex in the warm gas phase towards ζ Ophiuchi (Savage & Sembach 1996). Therefore, the $[O/Zn]$ ratio is an excellent dust-free diagnostic tool for the $[\alpha/\text{iron-peak}]$ abundance ratio.

From the average of the two transitions we derive $[O/H]=-1.86$, which implies $[O/Zn]=0.21$, a relatively modest, if any, α enhancement. Si abundance is almost the same as oxygen and we obtain $[Si/H]=-1.91$, which corresponds to $[Si/Zn]=0.16$. We note also here that Lu et al. (1996) tentatively identified the SII 1253.811 Å line associated with the damped, and derived a column density $\log N(\text{SII}) = 14.70 \pm 0.03$, i.e. an abundance of $[S/H]=-1.91$, which implies $[S/Zn]=0.16$. Thus, the S abundance of Lu et al. (1996) is also consistent with our O and Si abundances in the system.

For the first time in the DLA studied here we have at our disposal a set of 3 indicators for the α element abundances and three indicators for the iron peak elemental abundances, which do not need any dust correction. The $[O, Si, S/Zn, Fe, Cr]$ ratios, in whatever combination the elements are chosen, are in the range 0.09–0.22 dex. This range of values is significantly lower than analogous ratios in Galactic stars with comparable metallicities, and the difference is more stringent in this case for the low value of the metallicity, which is one order of magnitude lower than the average value in DLAs. The lack of significant $[\alpha/Fe]$ enhancement corroborates previous results on this system as well as the suggestion that, on the basis of their relative elements, at least some DLA galaxies seem to undergo a chemical evolution which differs from that of the Milky-Way. Centurión et al. (2000) showed that the $[S/Zn]$ ratios in DLAs are consistent with a decreasing trend with increasing metallicity. These ratios show also a correlation, although less clear-cut, with cosmic time, where the lower $[S/Zn]$ values occur mostly at low redshifts. The $[S/Zn]=0.16$ and $[O/Zn]=0.19$ for our DLA add a new point which supports the presence of such trends. Considering the extant data of $[S/Zn]$ and the value $[O/Zn]=0.20$ we derived here, the regression analysis yields the correlation $[S, O/Zn] = -0.36(\pm 0.11)[Zn/H] - 0.52$. The $\alpha/\text{iron-peak}$ ratio becomes solar at $[Fe/H] \approx -1.4$ and further decreases at higher metallicities.

Chemical evolutionary models predict solar $[\alpha/\text{Fe}]$ at low metallicities, when star formation proceeds in bursts separated by relatively long quiescent periods or whenever star formation is slower than that of our Galaxy. These conditions are met in dwarf galaxies, LSB galaxies and in the outer regions of disks (Matteucci et al. 1997, Jiménez et al. 1999). In these galaxies the metal enrichment is so slow that Type Ia supernovae have enough time to evolve and enrich the gas with iron-peak elements, in such a way that reduced α over iron ratios are attained at low metallicity.

In the Milky Way few metal poor stars with solar $[\alpha/\text{Fe-peak}]$ ratios have been found (Carney et al. 1997, Nissen & Schuster 1997). These exceptional stars show large apogalactic distances and it was suggested that they belonged to a satellite galaxy which experienced a different chemical evolution history than the Milky Way, and were subsequently accreted by the latter.

Incidentally, we note that the interpretation of the low $[\text{O}/\text{Fe}]$ ratio at low metallicities as an intrinsic effect of Type I SNe does not support the suggestion by Umeda et al. (1999) that Type I SNe may be suppressed at metallicities lower than $[\text{Fe}/\text{H}] < -1.0$.

4.3. The Nitrogen abundance

In our system the nitrogen abundances relative to Zn and O are $[\text{N}/\text{Zn}] = -0.54$ and $[\text{N}/\text{O}] = -0.74$. Both values are somewhat lower than previously reported by Molaro et al. (1996) because of the upwards revision in oxygen and iron abundances. The $[\text{N}/\text{O}]$ is not so extreme as previously thought, but the N abundance still requires a primary production of the element according to the evolutionary models discussed in Matteucci et al (1997). N abundance in this DLA, but also in other DLAs, remain lower than the corresponding values in the Milky Way, where N is found to trace Fe in lockstep (Israelian et al. 2000), which again suggests a different pattern than the Milky Way. Nitrogen abundances have been discussed by Lu, Sargent, & Barlow (1998) and by Centurión et al. (1998), who pointed out the presence of a real scatter among DLA systems with some values very close to a pure secondary behavior and other values which require a primary nucleosynthesis. The different values of N/O observed at a given O/H may be understood in terms of the delayed delivery of primary N with respect to O when star formation proceeds in bursts. During the quiescence period N is deposited in the ISM and the N/O ratio increases while O/H remains constant. Therefore a relatively high $[\text{N}/\text{O}]$, such as that observed here, nicely fits with the picture of a delayed N release after a quiescence period, which is also required by the low α over iron ratios.

4.4. Conclusions

UVES observations of the QSO 0000-2620 resulted in the detection of the ZnII 2026.136 Å transition originated in the damped system at $z_{abs}=3.3901$. The abundance derived is $[Zn/H]=-2.06$ which is presently the lowest among DLAs. This low metallicity level shows that the galaxy is in the early stages of its chemical evolution. When compared with the larger abundances observed in DLA galaxies at lower redshift, this measurement provides a first hint of a cosmological chemical evolution in which abundances increase with decreasing redshift (see Vladilo et al 2000)

The abundances of Cr and Fe were also obtained and found similar to the abundance of Zn. This coincidence in the abundances between refractory and non refractory elements is interpreted as dust has not formed yet.

The OI 925 Å and 950 Å lines were detected although they fall within the Lyman α forest. These lines have oscillator strengths much smaller than that of the OI 1302 Å line which is strongly saturated. These detections allow, for the first time, a rather accurate measurement of the oxygen abundance in a DLA system. The oxygen abundance, $[O/H]=-1.85$, is remarkably similar to the sulphur abundance derived by Lu et al (1996), $[S/H]=-1.91$, and both abundances are similar to that of silicon, $[Si/H]=-1.91$, which is consistent with the absence of dust in the system as implied by Zn and Fe abundances.

In this DLA the relative abundances of α and iron-peak elements are mildly overabundant compared to solar and do not show the enhancement expected for a progenitor of an early type spiral such as our own Galaxy. The paradigm of DLAs as progenitors of the Milky-Way type spirals is not supported by the study of the dust-independent chemical evolution indicators such as $[O/Zn]$ and $[S/Zn]$. The almost solar abundance ratios measured at the low metallicity of $[Fe/H]\approx -2$ suggest that this DLA should be associated with objects with low, or episodic, star formation rates such as LSB or dwarf galaxies. However, these results are not characteristic of this DLA alone and similar low $[S/Zn]$ ratios are also found in other DLAs (Centurión et al 2000).

5. Acknowledgements

The new high resolution spectra analyzed in this paper are of unique quality and were obtained during the first nights of commissioning of a new instrument at a new telescope. For these results we are indebted to all ESO staff involved in the VLT construction and in UVES commissioning.

REFERENCES

- Bergeson, S. D., Lawler J. E. 1993a ApJ408, 382
- Bergeson, S. D., Lawler J. E. 1993b ApJ414, 137
- Boesgaard A. M. King J. R., Deliyannis C. P., Vogt S. S. 1999 Astron J 117, 492
- Cardelli, J. A., & Savage B., D., 1995, ApJ452, 275
- Carney, B.W., Wright, J.S., Sneden, C., Laird, J.B., Aguilar, L.A., Latham, D.W. 1997, AJ114, 363
- Centurión, M., Bonifacio, P., Molaro, P., Vladilo, G. 1998, ApJ, 509, 620
- Centurión, M., Bonifacio, P., Molaro, P., Vladilo, G. 2000, ApJ, 536 in press astro-ph 0001109
- Dekker, H., D’Odorico, S. & Kaufer A., 2000, Proceedings of the SPIE Conference 4013, in press
- Fedchak J. A., Lawler J. E. 1999, ApJ523, 734
- Fontana, A. & Ballester, P. 1995, The Messenger, 80,37
- Fritze- v. Alvensleben, U., Lindner, U., Moller, C.S., and Fricke, K., J., 2000 astro-ph/9904208
- Grevesse, N. Noels A., Sauval A. J. 1996 in Cosmic Abundances ASP Conf Series Vol 99, 117
- Haehnelt, M.G., Steinmetz, M. & Rauch, M. 1998, ApJ, 495, 647
- Israelian G., Garcia Lopez, R., Rebolo R., 1998 ApJ507, 805
- Israelian G., Garcia Lopez R., Rebolo R., 2000 astro-ph/0001426.
- Jimenez, R., Bowen, D.V., & Matteucci, F. 1999, ApJ, 514, 83
- Ledoux, C., Petitjean, P., Bergeron, J., Wampler, E.J., & Srianand R. 1998, A&A, 337, 51
- Le Brun, V., Bergeron, J., Boisse, P., & Deharveng, J.M. 1997, A&A, 321, 733
- Lu, L., Sargent, W.L.W., Barlow T.A. & Churchill C.W. 1996, ApJS, 107,475
- Lu, L., Sargent, W.L.W., & Barlow, T.A. 1998, AJ, 115, 55

- Lu, L., Sargent, W.L.W., Barlow, T.A., Churchill, C.W., Vogt, S. 1996, ApJS, 107, 475
- Matteucci, F., Molaro, P., & Vladilo, G. 1997, A&A, 321, 45
- Molaro, P., Centuri3n, M., & Vladilo, G. 1998, MNRAS, 293, L37
- Molaro, P., D’Odorico, S., Fontana, A., Savaglio, S., Vladilo, G. 1996, A&A308, 1
- Morton, D.C. 1991, ApJS, 77, 119
- Nissen, P.E., & Schuster, W.J. 1997 A&A, 1997, 326, 751
- Outram P. J., Chaffee F. H., and Carswell R. F. 1999 MNRAS310, 289
- Pettini, M., King, D.L., Smith, L.J., Hunstead, R.W. 1997b, ApJ, 478, 536
- Pettini, M., Smith, L.J., Hunstead, R.W., King, D.L. 1994, ApJ, 426, 79
- Pettini, M., Ellison, S.L., Steidel, C.C., Bowen, D.V. 1999, ApJ, 510, 576
- Pettini M., Ellison S., Steidel C., Shapley A. E., Bowen D. V. 2000, ApJin press (astro-ph/9910131)
- Pettini, M., Smith, L.J., King, D.L., Hunstead, R.W. 1997, ApJ, 486, 665
- Prantzos N., Boisser S. 2000 MNRASin press (astro-ph/0001313)
- Prochaska, J.X. & Wolfe, A.M. 1997, ApJ, 487, 73
- Prochaska, J.X. & Wolfe, A.M. 1999, ApJS, 121, 369
- Rao, S.M., & Turnshek, D.A. 1998, ApJ, 500, L115
- Rao, S.M., & Turnshek, D.A. 2000, astro-ph 9909164
- Savage, B.D., & Sembach, K.R. 1996, Ann. Rev. Astron. Astrophys., 34, 279
- Savaglio S. Panagia, N., Stiavelli, M., 2000 in *Cosmic Evolution and Galaxy Formation* ASP Conf Series, eds J. Franco, E. Terlevich, O. L3pez-Cruz, I. Aretxaga in press (astro-ph/9912112)
- Umeda H., Nomoto K., Kobayashi C., Hachisu I., Kato M., 1999 ApJ522 L43
- Viegas S. M. 1995 MNRAS276, 268

Vladilo, G., 1998, ApJ, 493, 583

Vladilo, G., 1999, in *The Evolution of Galaxies on Cosmological Timescales* ASP Conference Series, eds. J.E. Beckman, and T.J. Mahoney (astro-ph/9903406).

Vladilo, G. Bonifacio, P., Centuri3n, M., Molaro, P., 2000 ApJsubmitted.

Wolfe, A.M., Lanzetta, K.M., Foltz, C.B., Chaffee, F.H. 1995, ApJ, 454, 698

Figure caption:

Fig. 1.— Observed lines for the α -elements O and Si associated with the $z_{abs}=3.3901$ DLA system towards QSO 0000–2620. All the lines are aligned taking zero velocity in correspondence of $z_{abs}=3.39010$. Smooth lines are the synthetic spectra obtained from the fit as described in the text. The dotted line in the bottom panel bluewards the SiII feature is the CIV $\lambda\lambda$ 1548 Å of the $z_{abs}\approx 4.126$ metallic system. The absorption features in the red side are the CIV $\lambda\lambda$ 1548 Å of the $z_{abs}\approx 4.130$ and 4.131. Cosmic removals are marked by crosses.

Fig. 2.— As Fig 1 for the nitrogen lines. The top panel is centered on the NI $\lambda\lambda$ 1134.4149 Å transition, with the other two lines NI $\lambda\lambda$ 1134.1653 Å and NI $\lambda\lambda$ 1134.9803 Å on the blue and red sides, respectively.

Fig. 3.— As Fig 1 for the iron-peak elements. The two dotted features on both sides of CrII $\lambda\lambda$ 2056.2539 Å are telluric lines.

Table 1. Column densities.

Ion	λ	f ^a	log (N)	b	z_{abs}	(Z/H) _⊙	[Z/H]	[Z/Zn]
HI	1215.670	0.4164	21.41 ± 0.08 ^b					
NI	953.6549	0.02032	14.77±0.03	9.4±0.7	3.390163	−4.03±0.07	−2.61	−0.54
NI ^f	1134.1653	0.01342	14.70± 0.02	10.0 ± 0.1	3.390123	−4.03±0.07	−2.68	−0.61
NI ^f	1134.4149	0.02683						
NI ^f	1134.9803	0.04023						
OI	925.4460	0.0003539	16.50 ± 0.03	14.3 ± 1.5	3.390161	−3.13 ± 0.07	−1.78	0.29
OI	950.8846	0.001571	16.35±0.08	10.9± 0.9	3.390080	−3.13 ± 0.07	−1.93	0.14
FeII	1611.2004	0.001020 ^c	14.87 ± 0.03	9.7 ± 1.3	3.390105	−4.50 ± 0.01	−2.04	0.03
SiII	1808.0126 ^f	0.002180	15.06±0.02	9.8 ± 0.3	3.390102	−4.44±0.01	−1.91	0.16
CrII ^f	2056.2539	0.1052 ^d	13.09 ± 0.03	8.8 ± 0.9	3.390078	−6.31 ± 0.01	−2.01	0.06
CrII ^f	2062.2339	0.07796 ^d						
ZnII	2026.1360	0.4886 ^d	12.01±0.05	9.3±1.3	3.390144	−7.33±0.04	−2.07	—
NiII	1709.600	0.03482 ^e	13.39 ± 0.03	8.99 ± 0.80	3.390127	−5.75 ± 0.01	−2.27	−0.20

REFERENCES— (a) Unless otherwise indicated the f values are from Morton (1991); (b) Prochaska & Wolfe (1999); (d) Bergeson & Lawler (1993a); (c) Cardelli & Savage (1995); (e) Fedchak & Lawler (1998); (f) Bergeson & Lawler (1993b)

NOTES—(f) The N λ 1134 Å was treated as a multiplet in the fitting procedure; (g) All abundances were normalised to the meteorite, when available, or to photospheric values reported by Grevesse et al. (1996).





